


# Gadobutrol

## Gd-DO3A-butrol

Created: February 15, 2006

Updated: March 13, 2006

<b>Chemical name:</b>	Gadobutrol
<b>Abbreviated name:</b>	Gd-DO3A-butrol, Gd-BT-DO3A
<b>Synonym:</b>	[1,4,7-Tris(carboxymethyl)-10-(1-(hydroxymethyl)-2,3-dihydroxypropyl)-1,4,7,10-tetraazacyclododecanato] gadolinium(III); Gadovist®; 10-[(1 <i>SR</i> ,2 <i>RS</i> )-2,3-dihydroxy-1-hydroxymethylpropyl]-1,4,7,10-tetraazacyclodecane-1,4,7-triacetic acid, gadolinium
<b>Backbone:</b>	Compound
<b>Target:</b>	Blood pool, extracellular fluid space
<b>Mechanism:</b>	Nonspecific filling of extracellular fluid space
<b>Method of detection:</b>	MRI
<b>Source of signal:</b>	Gadolinium
<b>Activation:</b>	No
<b><i>In vitro</i> studies:</b>	Yes
<b>Rodent studies:</b>	Yes
<b>Other non-primate mammal studies:</b>	Yes
<b>Non-human primate studies:</b>	No
<b>Human studies:</b>	Yes



Click on the above structure for additional information in PubChem  
[<http://pubchem.ncbi.nlm.nih.gov>].

## Background

[PubMed]

Gadobutrol (Gd-DO3A-butrol) is a nonionic, paramagnetic contrast agent developed for tissue contrast enhancement in magnetic resonance imaging (MRI) (1-3).

Paramagnetic contrast agents are generally metal chelates with unpaired electrons, and they work by shortening both the  $t_1$  and  $t_2$  relaxation times of surrounding water protons to produce the signal-enhancing effect (1, 2). At normal clinical doses of 0.1-0.2 mmol/kg, the  $t_1$  effect tends to dominate. Current agents are water-soluble compounds that distribute in the extracellular fluid and do not cross the intact blood-brain barrier (BBB). They are often used to enhance signals of central nervous system (CNS) tissues that lack a BBB (e.g., pituitary gland), extraaxial tumors (e.g., menin-

giomas), and areas of BBB breakdown (e.g., tumor margins). In these cases, small or multiple CNS lesions are more clearly delineated with contrast enhancement. In addition, contrast enhancement can highlight vasculature, delineate the extent of disease, and confirm the impression of normal or nonmalignant tissues. These contrast agents can also be used in a similar nonspecific manner to enhance contrast between pathologies and surrounding normal areas in other organs (1, 2, 4).

Gadolinium(III) ( $\text{Gd}^{3+}$ ), a lanthanide metal ion with seven unpaired electrons, has been shown to be very effective at enhancing proton relaxation because of its high magnetic moment and very labile water coordination (2, 5-7). Gadopentetate dimeglumine (Gd-DTPA) was the first intravenous MRI contrast agent used clinically, and a number of similar gadolinium chelates have been developed in an effort to further improve clinical efficacy, patient safety and patient tolerance. The major chemical differences among these Gd chelates are the presence or absence of overall charge, ionic or nonionic, and their ligand frameworks (linear or macrocyclic). Gd-DO3A-butrol has a macrocyclic framework and is neutral. The DO3A-butrol ligand was developed based on the belief that high overall hydrophilicity of an agent is generally associated with very low protein binding and good biological tolerance (8). It is a modification of the Gd-DO3A compound in which a trihydroxybutyl group is attached to the macrocyclic ligand to ensure high hydrophilicity. The ligand contains two chiral carbons at C-13 and C-14, therefore, the product is a racemic mixture of (13*R*,14*S*)– and (13*S*,14*R*)–. The Gd(III) in Gd-DO3A-butrol has a coordination number of 9. Gd-DO3A-butrol is a water-soluble, highly hydrophilic compound with a partition coefficient between n-butanol and buffer at pH 7.6 of ~ 0.006 (9).

Gd-DO3A-butrol is not commercially available in the United States, but it is commercially available in Canada at a concentration of 1.0 mmol/ml (604.72 mg/ml) for contrast enhancement during cranial and spinal imaging with MRI and MR angiography. At this concentration, it has an osmolarity of 1117 mOsm/kg at 37 °C, viscosity of 4.96 mPa at 37 °C, and log *P*(n-octanol/water) of –5.4 at 25 °C.

## Synthesis

---

[PubMed]

Vogler et al. (10) discussed the general method for preparing the macrocyclic core based on the Richman-Atkins method. In this method, 5-amino-2,2-dimethyl-6-hydroxy-1,3-dioxepan was treated with tosyl aziridine to give 1,7-ditoxyl-4-(6-hydroxy-2,2-dimethyl-1,3-dioxepan-5-yl)-1,4,7-triazaheptane. This compound was deprotonated with sodium ethoxide and reacted with the tritosylate of diethanolamine to give a macrocyclic intermediate. It was then treated with lithium in liquid ammonia to remove the tosyl groups to give a mono-substituted tetramine, which was then alkylated with chloroacetic acid at pH 9-10. DO3A-butrol was produced by a final treatment with diluted hydrochloric acid. The ligand was reacted with gadolinium oxide in water at 90 °C to produce Gd-DO3A-butrol.

Platzek et al. (8) described three approaches to synthesizing the DO3A-butrol ligand. One approach was preferred because it involved fewer steps and allowed scaled up production for clinical applications. In this approach, 1-(1-(hydroxymethyl)-2,3-dihydroxypropyl)-1,4,7,10-tetraazacyclododecane tetrahydrochloride was first prepared. The compound was then reacted with

chloroacetic acid at 70 °C and pH 9-10 for 18 h. The yield was 65%. Gd-DO3A-butrol was then prepared by adding Gd<sub>2</sub>O<sub>3</sub> to the ligand in water and stirring at 90 °C for 6 h. Cation- and anion exchange resins were then added at room temperature, and the mixture was stirred for 30 min. The resins were recollected by filtration, and charcoal was added to the filtrate and refluxed for 1 h. The solution was filtered and evaporated to dryness. The residue was redissolved in water and ethanol, and then was refluxed for 2 h. Gd-DO3A-butrol was collected by filtration and drying under vacuum (12 h at 60 °C). The yield was 87%. Mass spectroscopy (FAB) gave the following results: *m/e* 606 [(M + H)<sup>+</sup>]; *m/e* 1211 [(2M + H)<sup>+</sup>]. The infrared spectrum (KBr) had peaks at 3560, 3280, 2980, 2975, 2940, 2920, 2880, 2870, 1650, 1600, and 1380 cm<sup>-1</sup>. The structure was determined by X-ray crystallography.

## ***In Vitro* Studies: Testing in Cells and Tissues**

---

[PubMed]

Pintaske et al. (11) determined the *in vitro* relaxivity of Gd-DO3A-butrol and other compounds (Gd-DTPA and Gd-BOPTA) in human plasma obtained from healthy volunteers. Measurements of Gd-DO3A-butrol in concentrations of 0.01 mM up to 64 mM at 37 °C were made with 0.2-, 1.5-, and 3.0- Tesla (T) clinical scanners. The longitudinal relaxation rates (*r*<sub>1</sub>; liter x mmol<sup>-1</sup> x s<sup>-1</sup>) of Gd-DO3A-butrol were 5.5 ± 0.3, 4.7 ± 0.2, and 3.6 ± 0.2, at 0.2, 1.5, and 3T, respectively. The transverse relaxation rates (*r*<sub>2</sub>; liter x mmol<sup>-1</sup> x s<sup>-1</sup>) of were 10.1 ± 0.3, 6.8 ± 0.2, and 6.3 ± 0.3, at 0.2, 1.5, and 3 T, respectively. In comparison, these values were lower for Gd-DTPA but higher for Gd-BOPTA.

Vogler et al. (10) measured the *in vitro* physico-chemical properties of Gd-DO3A-butrol. The osmolality was found to be 0.57 osmol/kg for the 0.5 M concentration and 1.39 osmol/kg for the 1 M concentration. The distribution coefficient (butanol/water) was 0.006, and the viscosity was 1.43 cP for the 0.5 M concentration and 3.7 cP for the 1 M concentration. The relaxivity (*t*<sub>1</sub>; liter x mmol<sup>-1</sup> x s<sup>-1</sup>) in plasma was determined to be 5.6 at 0.45 T and 6.1 at 2 T. There was negligible protein binding of about 2.7% in plasma. With use of rat mast cells, the *I*<sub>50</sub> (the concentration at which histamine was released) was found to be greater than 250 mM. The *I*<sub>50</sub> value for lysozyme inhibition (the concentration at which 50% of the enzyme was inhibited) was greater than 300 mM.

## **Animal Studies**

---

### **Rodents**

[PubMed]

Vogler et al. (10) studied the long-term elimination and biodistribution of 0.25 mmol/kg Gd-DO3A-butrol in rats. More than 90% of the i.p. injected dose (ID) was excreted by the kidneys in 2 h and 100% in 7 days. The total amount remaining in the body decreased from 0.71% ID on day 1 to 0.15% ID on day 7 after injection. Inductively-coupled plasma atomic emission spectrometry (detection limit was 0.01 ppm or 65 nM) showed that only the kidneys, bones, and blood contained an amount of the agent above the detection limit after 7 days. The % ID of these organs were 0.06

$\pm 0.01$ ,  $0.01 \pm 0.00$ , and  $0.01 \pm 0.00$ , respectively. The acute toxicity (lethal dose ( $LD_{50}$ )) in mice was 23 mmol/kg. In a neural tolerance study, the median  $LD_{50}$  (lethal dose;  $n = 10$ ) and effective dose ( $ED_{50}$ ) of Gd-DO3A-butrol in rats after intracisternal injection were 86  $\mu\text{mol/kg}$  and 18  $\mu\text{mol/kg}$ , respectively. In comparison, the  $LD_{50}$  and  $ED_{50}$  for Gd-DTPA were 740  $\mu\text{mol/kg}$  and 73  $\mu\text{mol/kg}$ , respectively. The study suggested that macrocyclic compounds had lower cerebral tolerance and that linear compounds had higher cerebral tolerance. Imaging studies in rat models with cerebral infarct, brain tumors, and intramuscular tumors showed contrast enhancements (0.3 mmol/kg Gd-DO3A-butrol) in all three pathologic areas. A dose-dependent increase in the signal intensity of the intramuscular tumors was observed.

Several rat studies (12-16) showed that Gd-DO3A-butrol was effective in providing contrast enhancement for MRI in glioma, brain ischemia, hepatocellular carcinoma, and the lymphatic system.

## Other Non-Primate Mammals

[PubMed]

Vogler et al. (10) performed Gd-DO3A-butrol pharmacokinetics studies in beagle dogs. The elimination  $t_{1/2}$  and the plasma clearance were  $45 \pm 3.6$  min and  $3.75 \pm 0.30$  ml/min/kg, respectively. The total volume of distribution at steady state ( $V_{ss}$ ) was  $0.23 \pm 0.02$  liter/kg. High-performance liquid chromatography analysis of the plasma and urine samples showed that Gd-DO3A-butrol was not metabolized.

The abdominal MRI enhancement properties of Gd-DO3A-butrol were investigated in pigs (17). Images of 40 and 80  $\mu\text{mol/kg}$  doses were conducted on a 1.5 T imager. Gd-DO3A-butrol gave signal intensity curves similar to those produced by extracellular agents. Significant enhancement was found in all tissues except the trunk muscles. Kalinoski et al. (18) used 1 M Gd-DO3A-butrol in a dose of 2 mmol/kg to demonstrate the feasibility of producing X-ray contrast enhancement with computed tomography (CT) in pigs

## Non-Human Primates

[PubMed]

## Human Studies

---

[PubMed]

Staks et al. (19) investigated the pharmacokinetics, dose proportionality, and tolerability of Gd-DO3A-butrol in healthy volunteers. Two double-blind, randomized, placebo-controlled phase I studies were conducted for testing a low concentration (0.5 M; doses = 0.04, 0.1, 0.2, 0.3, and 0.4 mmol/kg;  $n = 55$ ) and a high concentration (1 M; doses = 0.3, 0.4, and 0.5 mmol/kg;  $n = 36$ ). From the data for 24 volunteers, pharmacokinetic parameters were calculated from an open two-compartment model. The values for plasma elimination half-life ( $t_{1/2\beta}$ ),  $V_{ss}$  (liter/kg), renal clearance (ml/min/kg), and 72-h total renal excretion (Urine, 72 h (% ID)) for a 0.1 mmol/kg dose were  $1.78 \pm 0.43$ ,  $0.21 \pm 0.02$ ,  $1.56 \pm 0.18$ , and  $98 \pm 3.33$ , respectively. For a 0.4 mmol/kg dose, these values were  $1.33 \pm 0.21$ ,  $0.15 \pm 0.02$ ,  $1.45 \pm 0.22$ , and  $96.6 \pm 5.61$ , respectively. Plasma Gd-DO3A-butrol

decreased bioexponentially, and the volume of distribution was within the extracellular fluid space. Elimination occurred primarily through the renal route. Biotransformation analysis of the urine and plasma samples revealed no metabolites from Gd-DO3A-butrol. No significant changes were observed in serum chemistry/hematology, or urine chemistry/urinalysis. No clear dose-dependent adverse events were found between the low-dose and high-dose studies. The most common adverse events were smell sensation (12.5%) and taste sensation (15%).

Tombach and Heindel (20) and Huppertz and Rohrer (21) summarized clinical studies for the clinical applications of 1 M Gd-DO3A-butrol. In 29 phase I to phase III studies (2523 subjects and 2662 doses), Gd-DO3A-butrol was distributed in the extracellular fluid with a plasma terminal  $t_{1/2}$  of 1.7-2 h. Approximately 98% of the dose was excreted unchanged in the urine within 12 h. The studies indicated that 1 M Gd-DO3A-butrol had an excellent safety profile with doses ranging from 0.04 to 1.5 mmol/kg. Overall, 8.5% of the patients reported related and unrelated adverse events. For contrast enhancement, analysis of efficacy data showed results comparable to results to other commercially available 0.5 M Gd- chelates in studies of the CNS (677 patients), and angiography (676 patients).

In a study of the pharmacokinetics of 1 M Gd-DO3A-butrol in 21 patients with chronic renal failure, Tombach et al. (22) reported that the extracellular distribution of Gd-DO3A-butrol remained unchanged but the mean elimination  $t_{1/2}$  increased compared with that of healthy volunteers. Other human studies have shown the potential clinical usefulness of 1 M Gd-DO3A-butrol in various CNS and vascular diseases [PubMed].

## References

---

1. Brasch RC, Ogan MD, Engelstad BL. Paramagnetic Contrast Agents and Their Application in NMR Imaging, in Contrast media; Biologic effects and clinical application, Z. Parvez, R., R. Monada and M. Sovak, Editor. 1987, CRC Press, Inc.: Boca Raton, Florida. p. 131-143.
2. Saini S, Ferrucci JT. Enhanced Agents for Magnetic Resonance Imaging: Clinical Applications, in Pharmaceuticals in Medical Imaging, D.P. Swanson, H.M. Chilton and J.H. Thrall, Editor. 1990, MacMillan Publishing Co., Inc.: New York. p. 662-681.
3. multihance (gadobenate dimeglumine injection, 529 mg/ml) Package Insert. 2005: Bracco Diagnostics Inc. F1/3.541037.
4. Runge VM, Kirsch JE, Wells JW, Awh MH, Bittner DF, Woolfolk CE. Enhanced liver MR: Contrast agents and imaging strategy. *Crit Rev Diagn Imaging* 34(2):1-3; 1993. (PubMed)
5. DeSimone D, Morris M, Rhoda C, Lucas T, Zielonka J, Olukotun A, Carvlin M. Evaluation of the safety and efficacy of gadoteridol injection (a low osmolal magnetic resonance contrast agent). Clinical trials report. *Invest Radiol* 26:S212-S216; 1991. (PubMed)
6. Tweedle MF. Physicochemical properties of gadoteridol and other magnetic resonance contrast agents. *Invest Radiol* 27:S2-S6; 1992. (PubMed)
7. Yuh WT, Engelken JD, Muhonen MG, Mayr NA, Fisher DJ, Ehrhardt JC. Experience with high-dose gadolinium MR imaging in the evaluation of brain metastases. *AJNR Am J Neuroradiol* 13(1):335-345; 1992. (PubMed)
8. Platzek J, Blaszkiewicz P, Gries H, Luger P, Michl G, Muller-Fahrnow A, Raduchel B, Sulzle D. Synthesis and Structure of a New Macrocyclic Polyhydroxylated Gadolinium Chelate Used as a Contrast Agent for Magnetic Resonance Imaging. *Inorg Chem* 36(26):6086-6093; 1997. (PubMed)
9. 60472 mg/ml (10 mmol/ml) 2004, Berlex Canada, Inc: Quebec, Canada p 1-10 2004.

10. Vogler H, Platzek J, Schuhmann-Giampieri G, Frenzel T, Weinmann HJ, Raduchel B, Press WR. Pre-clinical evaluation of gadobutrol: a new, neutral, extracellular contrast agent for magnetic resonance imaging. *Eur J Radiol* 21(1):1–10; 1995. (PubMed)
11. Pintaske J, Martirosian P, Graf H, Erb G, Lodemann KP, Claussen CD, Schick F. Relaxivity of Gadopentetate Dimeglumine (Magnevist), Gadobutrol (Gadovist), and Gadobenate Dimeglumine (MultiHance) in Human Blood Plasma at 0.2, 1.5, and 3 Tesla. *Invest Radiol* 41(3):213–221; 2006. (PubMed)
12. Fink C, Bock M, Kiessling F, Delorme S. Interstitial magnetic resonance lymphography with gadobutrol in rats: evaluation of contrast kinetics. *Invest Radiol* 37(12):655–662; 2002. (PubMed)
13. Frigeni V, Miragoli L, Grotti A, Lorusso V. Comparative study between gadobenate dimeglumine and gadobutrol in rats with brain ischemia: evaluation of somatosensory evoked potentials. *Invest Radiol* 36(10):561–572; 2001. (PubMed)
14. Kawai J, Takahashi M, Kato N, Takashima K, Miyazawa T. Comparison of a 1.0 molar and a 0.5 molar formulation of gadobutrol in dynamic MR imaging of the liver in rats with hepatocellular carcinoma. *Magn Reson Imaging* 17(1):91–97; 1999. (PubMed)
15. Le Duc G, Corde S, Charvet AM, Elleaume H, Farion R, Le Bas JF, Esteve F. In vivo measurement of gadolinium concentration in a rat glioma model by monochromatic quantitative computed tomography: comparison between gadopentetate dimeglumine and gadobutrol. *Invest Radiol* 39(7):385–393; 2004. (PubMed)
16. Reimer P, Allkemper T, Bremer C, Rummeny EJ, Spiegel HU, Balzer T, Peters PE. Assessment of reperfusion injury by means of MR contrast agents in rat liver. *J Magn Reson Imaging* 7(3):490–494; 1997. (PubMed)
17. Adam G, Neuerburg J, Spuntrup E, Muhler A, Scherer K, Gunther RW. Dynamic contrast-enhanced MR imaging of the upper abdomen: enhancement properties of gadobutrol, gadolinium-DTPA-polylysine, and gadolinium-DTPA-cascade-polymer. *Magn Reson Med* 32(5):622–628; 1994. (PubMed)
18. Kalinowski M, Kress O, Wels T, Alfke H, Klose KJ, Wagner HJ. 1-molar gadobutrol as a contrast agent for computed tomography: results from a comparative porcine study. *Invest Radiol* 38(4):193–199; 2003. (PubMed)
19. Staks T, Schuhmann-Giampieri G, Frenzel T, Weinmann HJ, Lange L, Platzek J. Pharmacokinetics, dose proportionality, and tolerability of gadobutrol after single intravenous injection in healthy volunteers. *Invest Radiol* 29(7):709–715; 1994. (PubMed)
20. Tombach B, Heindel W. Value of 1.0- M gadolinium chelates: review of preclinical and clinical data on gadobutrol. *Eur Radiol* 12(6):1550–1556; 2002. (PubMed)
21. Huppertz A, Rohrer M. Gadobutrol, a highly concentrated MR-imaging contrast agent: its physicochemical characteristics and the basis for its use in contrast-enhanced MR angiography and perfusion imaging. *Eur Radiol* 14:M12–M18; 2004. (PubMed)
22. Tombach B, Bremer C, Reimer P, Schaefer RM, Ebert W, Geens V, Heindel W. Pharmacokinetics of 1M gadobutrol in patients with chronic renal failure. *Invest Radiol* 35(1):35–40; 2000. (PubMed)